




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : C07D 277/36, 277/34, A61K 31/425, C07D 417/06, 417/10		A1	(11) International Publication Number: <b>WO 00/18748</b>
			(43) International Publication Date: 6 April 2000 (06.04.00)
(41) International Application Number: PCT/EP99/07250		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 30 September 1999 (30.09.99)		<p>Published With international search report.</p>	
(30) Priority Data: 98118538.2 30 September 1998 (30.09.98) EP			
(71) Applicant (for all designated States except US): ROCHE DIAGNOSTICS GMBH [DE/DE]; D-68298 Mannheim (DE).			
(72) Inventors; and (75) Inventors/Applicants (for US only): ESSWEIN, Angelika [DE/DE]; Birkenweg 4, D-64572 Büttelborn (DE). SCHAEFER, Wolfgang [DE/DE]; Tannhaeuserring 190, D-68199 Mannheim (DE). TSAKLAKIDIS, Christos [GR/DE]; Huegelstrasse 1/1, D-69469 Weinheim (DE). HONOLD, Konrad [DE/DE]; Suedstrasse 24, D-82377 Penzberg (DE). KALUZA, Klaus [DE/DE]; Hochfeldanger 3, D-83670 Bad Heilbrunn (DE). HOFFMANN, Eike [DE/DE]; Rathausstrasse 71, D-68519 Viernheim (DE).			
(74) Common Representative: ROCHE DIAGNOSTICS GMBH; Patent Dept., D-68298 Mannheim (DE).			

Title: RHODANINE DERIVATIVES FOR THE TREATMENT AND PREVENTION OF METABOLIC BONE DISORDERS



## (57) Abstract

The object of the present invention are compounds of general formula (I), in which m signifies a number between 0 and 8; q signifies a number between 0 and 8; X signifies the group CH<sub>2</sub> or C=S, whereby A signifies a single bond and m signifies 0 when X signifies CH<sub>2</sub>; A signifies a single or double bond; R<sub>1</sub>, R<sub>2</sub> signify hydrogen or lower alkyl, whereby R<sub>1</sub> and R<sub>2</sub> can be the same or different and, when m signifies 2-8, R<sub>1</sub> and R<sub>2</sub> in the group CR<sub>1</sub>=CR<sub>2</sub> can have various significances within the following sequence; R<sub>3</sub> signifies hydrogen or lower alkyl; Z signifies oxygen, sulphur; W signifies an optionally mono- or polysubstituted saturated or unsaturated mono-, bi- or tricycle which can contain one or more hetero atoms, as well as their physiologically compatible salts, esters, optically active forms, racemates, tautomers, as well as derivatives which can be metabolized *in vivo* to compounds of general formula (I), as well as the use of these compounds for the production of medicaments for the prophylaxis or therapy of metabolic bone disorders.

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

## Rhodanine derivatives for the treatment and prevention of metabolic bone disorders

5

The present invention is concerned with rhodanine derivatives for the treatment and prevention of metabolic bone disorders, a process for their manufacture as well as medicaments which contain these compounds.

10

In healthy persons the synthesis and degradation processes in bones is almost in equilibrium, i.e. the activity of the osteoblasts and osteoclasts is balanced. However, if this equilibrium is disturbed in favour of the osteoclasts and/or to the detriment of the osteoblasts, this leads to a reduction in the bone mass and to a negative change in the bone structure and function.

15

Hitherto, bone resorption inhibitors such as oestrogens, calcitonin and biphosphonates have primarily been used for the treatment of metabolic bone disorders. The use of these substances is, however, limited and also does not show the desired effect in all cases. Compounds which have a stimulating activity on bone synthesis and in addition contribute to an increase in an already reduced bone mass are accordingly of especial significance for the treatment of metabolic bone disorders.

20

Compounds having the rhodanine structural element are known as antidiabetics, cytostatics, inflammation inhibitors and for the treatment of cardiovascular illnesses, e.g. WO9305039, WO 9705875, EP 677517.

25

The parathyroid hormone (PTH), a hormone from the parathyroid gland, is the natural ligand of the receptor and an important regulator for the maintenance of the calcium level in the body. PTH can stimulate bone formation or bone resorption. In this, it acts as a regulatory hormone on a series of enzymes, inter alia, on adenylate cyclase (cAMP synthesis) and on ornithine decarboxylase. PTH mobilizes calcium from bones in the case of calcium deficiency, reduces calcium excretion from the kidneys and simultaneously improves the resorption of calcium from the intestine by an increased synthesis of 1,25-(OH)<sub>2</sub>D<sub>3</sub>. A normalization of the calcium level is achieved by the

35


CONFIRMATION COPY

action on these target organs. On the other hand, the incorporation of calcium in bones is stimulated in the case of an elevated calcium level. This osteoanabolic activity of PTH and its fragments has been attributed to the activation of adenylate cyclase and of cAMP-dependent protein kinases (Rixon, R. Whitfield, J. et al JMBR 2 (8) 1179-89 (1994)).

Surprisingly, it has now been found that rhodanine derivatives of the present invention stimulate the PTH receptor-mediated cAMP formation. Compounds of the present invention are accordingly suitable for the broad treatment of metabolic bone disorders. They can be used primarily to good effect where the bone synthesis is disturbed, i.e. they are especially suitable for the treatment of osteopenic disorders of the skeletal system such as e.g. osteoporosis, inter alia, osteogenesis imperfecta as well as for the local assistance in bone regeneration and osteoinduction such as e.g. in orthopedic and maxillary medical indications, in fracture healing, osteosyntheses, pseudoarthroses and for the healing in of bone implants. However, having regard to these properties they also find use in the prophylaxis of osteoporosis.

By their influence on bone metabolism medicaments with the rhodanine derivatives of the present invention as active substances furthermore form a basis for the local and systemic treatment of rheumatoid arthritis, osteoarthritis and degenerative arthrosis.

The object of the present invention are compounds of general formula (I),



in which

- m signifies a number between 0 and 8,
- q signifies a number between 0 and 8
- 30 X signifies the group CH<sub>2</sub> or C=S, whereby A signifies a single bond and m signifies 0 when X signifies CH<sub>2</sub>,
- A signifies a single or double bond

R<sub>1</sub>, R<sub>2</sub> signify hydrogen or lower alkyl, whereby R<sub>1</sub> and R<sub>2</sub> can be the same or different and, when m signifies 2-8, R<sub>1</sub> and R<sub>2</sub> in the group CR<sub>1</sub>=CR<sub>2</sub> can have various significances within the following sequence

R<sub>3</sub> signifies hydrogen or lower alkyl

5 Z signifies oxygen, sulphur

W signifies an optionally mono- or polysubstituted saturated or unsaturated mono- (sic), bi- or tricycle which can contain one or more hetero atoms,

10 As a rule, lower alkyl signifies linear or branched alkyl residues with one to six carbon atoms, preferably methyl, ethyl, propyl, i-propyl, butyl, t-butyl, pentyl, hexyl, particularly methyl.

Alkoxy groups signify a combination of a C<sub>1</sub>-C<sub>10</sub>-alkyl group in accordance with the  
15 above definition with an oxygen atom, e.g. methoxy, ethoxy, propoxy, isopropoxy, butoxy and pentoxy groups.

Under monocycle there are to be understood optionally mono- or polysubstituted saturated or unsaturated ring systems with 3-8, preferably 5-7 carbon atoms, which  
20 optionally can be interrupted by one or more hetero atoms, such as nitrogen, oxygen or sulphur, especially the phenyl, pyridyl, pyrimidinyl, pyridazinyl, pyrazinyl, cyclopentyl, cyclohexyl, cyclohexenyl, cycloheptyl, morpholinyl, thiamorpholinyl, piperidinyl, piperazinyl, tetrahydrofuranyl, tetrahydropyranyl, furyl, thiophenyl, imidazolyl, thiazolyl, oxazolyl, isothiazolyl, isoxazolyl, 1,2,3-triazolyl or 1,2,4-triazolyl residue, as  
25 well as residues such as e.g. phenyl phenyl ether, diphenylmethane and biphenyl. Substituents are preferably lower alkyl, alkoxy, alkoxycarbonyl, alkylcarbonyl, hydroxy, benzyl, benzyloxy, phenyl, dioxymethylene, cyanobenzoxymethyl, pyrrolidine, alkoxhydroxy, carboxyl, dialkylamino, styryl and halogen.

30 In the case of the bicycle set forth under W, this is preferably a residue such as the naphthyl, tetrahydronaphthyl, decaliny, quinolinyl, chromane, chromene, isoquinolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, indolyl, benzimidazolyl, indazolyl, oxindolyl, benzofuranyl, benzothiophenyl, benzothiazolyl, benzoxazolyl or purinyl residue, especially the indolyl, naphthyl, benzimidazolyl, quinolinyl,  
35 tetrahydroquinolinyl, benzothiophenyl and benzofuranyl residue, which optionally can

be mono- or polysubstituted. Substituents are preferably lower alkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, alkoxycarbonyl, alkylcarbonyl, hydroxy, benzyl, benzyloxy, phenyl, dioxymethylene, cyanobenzoxymethyl, pyrrolidine, alkoxyhydroxy, carboxyl, dialkylamino, styryl and halogen.

5

Tricycle signifies anthracene, fluorene, dibenzofuran, dibenzooxepine or carbazole.

Compounds of formula I, wherein W is phenyl, naphthyl, indolyl or thienyl, X<sub>n</sub>C = S, Z = oxygen and m and g are both 0, are disclosed in EP-A-0677517 and WO-A-96/26207,

10 however for the treatment of Alzheimer's disease or as hypoglycemic agents.


Compounds of formula I, wherein W is phenyl, furyl, thienyl or pyrrolyl, X is C = S, Z is oxygen, A is a double bond and m is 0 or 1 and g is unequal 0 or n is unequal 0 and g is 2 are disclosed in EP-A-0398179, however as aldose reductase inhibitor.

15

Compounds of formula I, wherein W is endolyl, X is C = S, Z is oxygen, A is a double bond and m is 1 and g is 0 is disclosed in WO-A-98/01445, however as ATP-ase inhibitors.

20 Compounds formula I, wherein W is 4-(2,5-di-tert. butyl-phenol) and X is methylene are disclosed in EP-A-0211670, however for the treatment of inflammations.

Therefore subject of the present invention are also new compounds of formula I



5

in which

m signifies a number between 0 and 8,

q signifies a number between 0 and 8

10 X signifies the group CH<sub>2</sub> or C=S, whereby A signifies a single bond and m signifies 0 when X signifies CH<sub>2</sub>,

A signifies a single or double bond

R<sub>1</sub>, R<sub>2</sub> signify hydrogen or lower alkyl, whereby R<sub>1</sub> and R<sub>2</sub> can be the same or different and, when m signifies 2-8, R<sub>1</sub> and R<sub>2</sub> in the group CR<sub>1</sub>=CR<sub>2</sub> can have various significances within the following sequence

15

R<sub>3</sub> signifies hydrogen or lower alkyl

Z signifies oxygen, sulphur

W signifies an optionally mono- or polysubstituted saturated or unsaturated mono- (sic), bi- or tricycle which can contain one or more hetero atoms,

20

whereas W is not phenyl, naphthyl, indolyl or thienyl, if X is C = X, Z is sulfur and m and g are both 0,

25

whereas W is not phenyl, furyl, thienyl or pyrrolyl, if XnC = S, Z is sulfur, A is a double bond and m is 0 or 1 and g is unequal 0 or m is unequal 0 and g is 2,

whereas W is not indolyl, if X is C = S, Z is sulfur, A is a double bond and m is 1 and g is 0,

30


whereas W is not 4-(2,5-di-tert. butyl-phenyl), if X is methylene, as well as their physiologically compatible salts, esters, optically active forms, racemates, tautomers, as

well as derivatives which can be metabolized *in vivo* to compounds of general formula (I), as well as the use of these compounds for the production of medicaments.

- 5 Preferred are compounds of general formula I in which X signifies C=S, Z signifies oxygen, A signifies a double bond, m signifies a number from 0 to 2, q signifies 0 or 1, R<sub>1</sub> and R<sub>2</sub> respectively signify hydrogen or methyl, R<sub>3</sub> signifies hydrogen or methyl and W signifies a phenyl, naphthyl, thiophenyl, benzothiophenyl, furanyl, phenyl, pyridyl, cyclohexenyl, dibenzooxepinyl, pyrrol or imidazolyl residue, which optionally can be  
 10 mono- or polysubstituted by halogen, hydroxy, methoxy, ethoxy, benzyloxy, butoxycarbonyl, methyl, i-propyl, t-butyl, dioxymethylene, cyanobenzoxymethyl or benzyl.


The manufacture of the compounds of general formula (I) is possible according to  
 15 methods known per se. An overview of the methods of synthesis is set forth in Scheme 1 (J. Med. Chem. 37 322-8 (1994); Chem. Pharm. Bull. 30 3563-73 (19982); Chem. Heterocycl. Compd. EN 2 267-70 (1996); J. Med. Chem. 21 82-7 (1978); J. Org. Chem. 57 4047-49 (1992); T.L. 35 6971-74 (1994)); R signifies the group:

20





Scheme 1



The  $\alpha$ -halocarboxylic acids and aldehydes used as starting materials are either commercially available, known or can be prepared analogously to the generally known processes.

Compounds of formula (I) can be administered (sic) in liquid, solid or aerosol form orally, enterally, parenterally, topically, nasally, pulmonary or rectally in all usual non-toxic pharmaceutically acceptable carrier materials, adjuvants and additives. The compounds of formula (I) can also be applied locally to/in the bones (optionally with surgical intervention). The term parenteral embraces subcutaneous, intravenous and intramuscular delivery or infusions. Oral administration forms can be e.g. tablets, capsules, dragees, syrups, solutions, suspensions, emulsions, elixirs etc., which can contain one or more additives from the following groups, such as flavourings, sweeteners, colouring agents and preservatives. Oral administration forms contain the active ingredient together with non-toxic, pharmaceutically acceptable carrier materials which are suitable for the production of tablets, capsules, dragees etc., such as e.g. calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate;

starch, mannitol, methylcellulose, talc, highly dispersible silicic acids, high molecular fatty acids (such as stearic acid), groundnut oil, olive oil, paraffin, miglyol, gelatine, agar-agar, magnesium stearate, beeswax, cetyl alcohol, lecithin, glycerol, animal and vegetable fats, solid high molecular polymers (such as polyethylene glycol). Tablets, capsules, dragees etc. can be provided with an appropriate coating, e.g. glyceryl mono-  
5 stearate or glyceryl distearate, in order to prevent undesired side effects in the gastrointestinal tract or to give a longer duration of action by the delayed absorption in the gastrointestinal tract. As the injection medium there are preferably used sterile injectable aqueous or oily solutions or suspensions which contain the usual additives  
10 such as stabilizers and solubilizers. Such additives can be e.g. water, isotonic saline, 1,3-butanediol, fatty acids (such as oleic acid), mono- and diglycerides or miglyol. For rectal use there can be used all suitable non-irritating additives which are solid at normal temperatures and liquid at rectal temperatures, such as e.g. cocoa butter and polyethylene glycol. Pharmaceutically usual carrier media are used for application as  
15 aerosols. Creams, tinctures, gels, solutions or suspensions etc. with the pharmaceutically usual additives are used for external application. The dosage can depend on a variety of factors such as mode of administration, species, age and/or individual condition. The doses to be administered daily or at intervals lie at 1-1000 mg/individual, preferably at 10-250 mg/individual, and can be taken at one time or  
20 divided over several times.

The compounds of formula (I) can also be applied locally to/in the bones (optionally with surgical intervention). The application directly to/in the bones (optionally with surgical intervention) can be effected locally or carrier-bonded either in solution or  
25 suspension, conveniently by infusion or injection. Carrier-bonded compounds of formula (I) can be administered, for example, as gels, pastes, solids or as a coating on implants.

Biocompatible and preferably biodegradable materials are used as the carrier.  
30 Preferably, the materials themselves also induce wound healing or osteogenesis.

For local application it is preferred that the compounds of formula (I) are imbedded in polymer gels or films in order to immobilize them and to apply these preparations directly on the site of the bone to be treated. Such polymer-based gels or films consist,  
35 for example, of glycerine, methylcellulose, hyaluronic acid, polyethylene oxides and/or

poloxamers. Also suitable are collagen, gelatines and alginates and are described, for example, in WO 93/00050 and WO 93/20859. Further polymers are polylactic acid (PLA) and copolymers of lactic acid and glycolic acid (PLPG) (Hollinger et al., J. Biomed. Mater. Res. 17 71-82 (1983)) as well as the bone derivative "Demineralized Bone Matrix" (DBM) (Guterman et al. Kollagen Rel. Res. 8 419-4319 (1988)). Also suitable are polymers as are used, for example, for the adsorption of TGF $\beta$  and which are described in EP-A 0 616 814 and EP-A-0 567 391 and synthetic bone matrices in accordance with WO 91/18558.

10 Likewise suitable as carriers for the compounds of formula (I) are materials which are usually used for the implantation of bone substitutes or otherwise of therapeutically active substances. Such carriers are based, for example, on calcium sulphate, tricalcium phosphate, hydroxylapatite (sic) and its biodegradable derivatives and polyanhydrides. Apart from these biodegradable carriers there are also suitable carriers which are not  
15 biodegradable, but which are biocompatible. Such carriers are, for example, sintered hydroxylapatite, bioglass, aluminates or other ceramic materials (e.g. calcium aluminium phosphate). These materials are preferably used in combination with the biodegradable materials, such as especially polylactic acid, hydroxylapatite, collagen or tricalcium phosphate. Further non-degradable carriers are described, for example, in  
20 US Patent 4,164,560.

It is especially preferred to use a carrier which liberates the compounds of formula (I) continuously at the target site. Especially suitable for this are e.g. "slow release pellets" from Innovative Research of America, Toledo, Ohio, USA. Pellets which release the  
25 compounds of formula (I) over several days, preferably up to 100 days with a daily dosage of 1-10 mg/kg per day, are especially preferred.

Preferred in the scope of the present invention are, apart from the compounds named in the Examples and compounds derivable by a combination of all of the significances of  
30 the substituents set forth in the claims, the following derivatives as well as their physiologically compatible salts, esters, optically active forms, racemates, tautomers as well as derivatives which can be metabolized *in vivo* to compounds of general formula (I), as well as the use of these compounds for the production of medicaments,

## Preferred Compounds (PC):

1. 5-(9*H*-Fluoren-2-ylmethylene)-2-thioxo-thiazolidin-4-one
- 5 2. 5-Phenanthren-9-ylmethylene-thiazolidine-2,4-dithione
3. 5-Anthracen-9-ylmethyl-2-thioxo-thiazolidin-4-one
4. 5-(5-Furan-2-yl-penta-2,4-dienylidene)-2-thioxo-thiazolidin-4-one
5. 5-(2-Methoxy-benzylidene)-thiazolidine-2,4-dithione
6. 5-(2,3-Dimethoxy-benzyl)-2-thioxo-thiazolidin-4-one
- 10 7. 5-[3-(2,4-Dimethoxy-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one
8. 2-Thioxo-5-(2,4,5-trimethoxy-benzylidene)-thiazolidin-4-one
9. 5-(2,4,6-Trimethoxy-benzylidene)-thiazolidine-2,4-dithione
10. 5-(2,5-Dimethoxy-benzyl)-2-thioxo-thiazolidin-4-one
11. 5-[3-(2-Hydroxy-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one
- 15 12. 5-(2-Hydroxy-3-methoxy-benzylidene)-2-thioxo-thiazolidin-4-one
13. 5-(3-Ethoxy-2-hydroxy-benzylidene)-thiazolidine-2,4-dithione
14. 5-(2,3-Dihydroxy-benzyl)-2-thioxo-thiazolidin-4-one
15. 5-[3-(4-Diethylamino-2-hydroxy-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one
16. 5-(2-Hydroxy-4-methoxy-benzylidene)-2-thioxo-thiazolidin-4-one
- 20 17. 5-(2,4,6-Trihydroxy-benzylidene)-thiazolidine-2,4-dithione
18. 5-(2-Hydroxy-5-methoxy-benzyl)-2-thioxo-thiazolidin-4-one
19. 2-Thioxo-5-(3-*o*-tolyl-allylidene)-thiazolidin-4-one
20. 5-(4-Methoxy-2,3-dimethyl-benzylidene)-2-thioxo-thiazolidin-4-one
21. 5-(2,4,6-Trimethyl-benzylidene)-thiazolidine-2,4-dithione
- 25 22. 5-(2,5-Dimethyl-benzyl)-2-thioxo-thiazolidin-4-one
23. 5-[3-[3-(4-Methoxy-phenoxy)-phenyl]-allylidene]-2-thioxo-thiazolidin-4-one
24. 5-[3-(4-*tert*-Butyl-phenoxy)-benzylidene]-2-thioxo-thiazolidin-4-one
25. 5-(3-*p*-Tolyloxy-benzylidene)-thiazolidine-2,4-dithione
26. 5-(3-Methoxy-benzyl)-2-thioxo-thiazolidin-4-one
- 30 27. 2-Thioxo-5-[3-(3,4,5-trimethoxy-phenyl)-allylidene]-thiazolidin-4-one
28. 5-(4-Benzylloxy-3-methoxy-benzylidene)-2-thioxo-thiazolidin-4-one
29. 5-(3,5-Dimethoxy-benzylidene)-thiazolidine-2,4-dithione
30. 5-(3-Benzylloxy-benzyl)-2-thioxo-thiazolidin-4-one
31. 5-[3-(3-Hydroxy-4-methoxy-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one
- 35 32. 5-(3,4-Dihydroxy-benzylidene)-2-thioxo-thiazolidin-4-one

33. 5-(3-Methyl-benzylidene)-thiazolidine-2,4-dithione  
34. 5-(4-Methoxy-3-methyl-benzyl)-2-thioxo-thiazolidin-4-one  
35. 5-[3-(4-Diethylamino-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one  
36. 5-(4-Phenoxy-benzylidene)-2-thioxo-thiazolidin-4-one  
5 37. 5-(4-Methoxy-benzylidene)-thiazolidine-2,4-dithione  
38. 5-(3-Benzoyloxy-4-methoxy-benzyl)-2-thioxo-thiazolidin-4-one  
39. 5-[3-(4-Ethoxy-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one  
40. 5-(4-Butoxy-benzylidene)-2-thioxo-thiazolidin-4-one  
41. 5-Naphthalen-1-ylmethylene-thiazolidine-2,4-dithione  
10 42. 5-(2-Methoxy-naphthalen-1-ylmethyl)-2-thioxo-thiazolidin-4-one  
43. 5-[3-(4-Methoxy-naphthalen-1-yl)-allylidene]-2-thioxo-thiazolidin-4-one  
44. 5-Naphthalen-2-ylmethylene-2-thioxo-thiazolidin-4-one  
45. 5-(3,4-Bis-benzoyloxy-benzylidene)-thiazolidine-2,4-dithione  
46. 5-(9-Ethyl-9H-carbazol-3-ylmethyl)-2-thioxo-thiazolidin-4-one  
15 47. 5-[3-(5-Methoxy-1H-indol-3-yl)-allylidene]-2-thioxo-thiazolidin-4-one  
48. 5-Benzo[1,3]dioxol-5-ylmethylene-2-thioxo-thiazolidin-4-one  
49. 5-Quinolin-4-ylmethylene-thiazolidine-2,4-dithione  
50. 5-(4-Hydroxy-benzyl)-2-thioxo-thiazolidin-4-one  
51. 5-[3-(4-Hydroxy-3,5-dimethoxy-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one  
20 52. 5-(3-Ethoxy-4-hydroxy-benzylidene)-2-thioxo-thiazolidin-4-one  
53. 5-(4-Hydroxy-3,5-dimethyl-benzylidene)-thiazolidine-2,4-dithione  
54. 5-Biphenyl-4-ylmethyl-2-thioxo-thiazolidin-4-one  
55. 5-[3-(4-Isopropyl-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one  
56. 5-(4-Methyl-benzylidene)-2-thioxo-thiazolidin-4-one  
25 57. 5-(4-Ethyl-benzylidene)-thiazolidine-2,4-dithione  
58. 5-(2,2-Diphenyl-ethyl)-2-thioxo-thiazolidin-4-one  
59. 5-(2-Pentyl-3-phenyl-allylidene)-2-thioxo-thiazolidin-4-one  
60. 5-(2-Hexyl-3-phenyl-allylidene)-thiazolidine-2,4-dithione  
61. 5-Phenethyl-2-thioxo-thiazolidin-4-one  
30 62. 5-(5-Phenyl-penta-2,4-dienylidene)-2-thioxo-thiazolidin-4-one  
63. 5-[3-(2-Methoxy-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one  
64. 5-[3-(4-Dimethylamino-phenyl)-allylidene]-thiazolidine-2,4-dithione  
65. 5-(3-Phenyl-propyl)-2-thioxo-thiazolidin-4-one  
66. 5-[3-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-allylidene]-2-thioxo-thiazolidin-4-one  
35 67. 5-(3-Ethoxy-4-methoxy-benzylidene)-2-thioxo-thiazolidin-4-one

68. 5-(4-Diethoxymethyl-benzylidene)-thiazolidine-2,4-dithione  
69. 5-(4-Dimethylamino-naphthalen-1-ylmethyl)-2-thioxo-thiazolidin-4-one  
70. 5-[3-(2,6-Dimethoxy-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one  
71. 5-(2,4-Dimethoxy-3-methyl-benzylidene)-2-thioxo-thiazolidin-4-one  
5 72. 5-(4-Styryl-benzylidene)-thiazolidine-2,4-dithione  
73. 5-[4-(3-Dimethylamino-propoxy)-benzyl]-2-thioxo-thiazolidin-4-one  
74. 5-[3-(2-Methyl-1*H*-indol-3-yl)-allylidene]-2-thioxo-thiazolidin-4-one  
75. 5-(4-Hydroxy-3-methyl-benzylidene)-2-thioxo-thiazolidin-4-one  
76. 5-(2-Allyloxy-benzylidene)-thiazolidine-2,4-dithione  
10 77. 5-(2-Hexyloxy-benzyl)-2-thioxo-thiazolidin-4-one  
78. 5-[3-(4-Propoxy-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one  
79. 5-(4-Pentyloxy-benzylidene)-2-thioxo-thiazolidin-4-one  
80. 5-(4-Octyloxy-benzylidene)-thiazolidine-2,4-dithione  
81. 5-(5-Benzylloxy-1*H*-indol-3-ylmethyl)-2-thioxo-thiazolidin-4-one  
15 82. 5-(3-Benzofuran-2-yl-allylidene)-2-thioxo-thiazolidin-4-one  
83. 5-(4-Pyrrolidin-1-yl-benzylidene)-2-thioxo-thiazolidin-4-one  
84. 5-(2,3,4,5,6-Pentamethyl-benzylidene)-thiazolidine-2,4-dithione  
85. 5-(2-Benzylloxy-benzyl)-2-thioxo-thiazolidin-4-one  
86. 5-[3-(3-Ethoxy-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one  
20 87. 5-(3,4-Dihydroxy-5-methoxy-benzylidene)-thiazolidine-2,4-dithione  
88. 5-(3,5-Dihydroxy-benzyl)-2-thioxo-thiazolidin-4-one  
89. 5-[3-(4-Ethoxy-3-methoxy-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one  
90. 5-(4-Hexyloxy-benzylidene)-2-thioxo-thiazolidin-4-one  
91. 5-(4-Heptyloxy-benzylidene)-thiazolidine-2,4-dithione  
25 92. 5-(7-Methoxy-benzo[1,3]dioxol-5-ylmethyl)-2-thioxo-thiazolidin-4-one  
93. 5-[5-(4-Methoxy-phenyl)-penta-2,4-dienylidene]-2-thioxo-thiazolidin-4-one  
94. 2-Thioxo-5-(2,4,5-trimethyl-benzylidene)-thiazolidin-4-one  
95. 5-(4-Decyloxy-benzyliden)-thiazolidine-2,4-dithione  
96. 5-[3-(2-*tert*-Butylsulphanyl-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one  
30 97. 5-(4-Butyl-benzylidene)-2-thioxo-thiazolidin-4-one  
98. 5-(2-Hydroxy-3-methyl-benzylidene)-thiazolidine-2,4-dithione  
99. 5-(4-*tert*-Butoxy-benzyl)-2-thioxo-thiazolidin-4-one  
100. 5-[3-(4-Hexyl-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one  
101. 5-(4-Octyl-benzylidene)-2-thioxo-thiazolidin-4-one  
35 102. 5-(4-Dodecyloxy-benzylidene)-thiazolidine-2,4-dithione

103. 5-(4-Pentyl-benzyl)-2-thioxo-thiazolidin-4-one  
104. 5-[3-(3-Amino-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one  
105. 5-(2-Ethoxy-naphthalen-1-ylmethylene)-2-thioxo-thiazolidin-4-one  
106. 5-(7-Methyl-1*H*-indol-3-ylmethylene)-thiazolidine-2,4-dithione  
5 107. 5-[3-(3,5-Dimethyl-1-phenyl-1*H*-pyrazol-4-yl)-allylidene]-2-thioxo-thiazolidin-4-one  
108. 5-(2,5-Dimethyl-1-phenyl-1*H*-pyrrol-3-ylmethylene)-2-thioxo-thiazolidin-4-one  
109. 5-[3-(2,2-Dimethyl-chroman-6-yl)-allylidene]-2-thioxo-thiazolidin-4-one  
110. 5-(4-Isopropoxy-benzylidene)-2-thioxo-thiazolidin-4-one  
10 111. 5-(4-Hydroxy-naphthalen-1-ylmethyl)-2-thioxo-thiazolidin-4-one  
112. 5-(5-Furan-2-yl-4-methyl-penta-2,4-dienylidene)-2-thioxo-thiazolidin-4-one  
113. 5-(2,3-Dihydro-benzofuran-5-ylmethylene)-thiazolidine-2,4-dithione  
114. 5-Quinolin-2-ylmethyl-2-thioxo-thiazolidin-4-one  
115. 5-[3-(4-Dibutylamino-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one  
15 116. 5-(4-Isobutyl-benzylidene)-2-thioxo-thiazolidin-4-one  
117. 5-[3-(4-Hydroxy-3-methoxy-phenyl)-allylidene]-thiazolidine-2,4-dithione  
118. 5-(6-Methoxy-naphthalen-2-ylmethyl)-2-thioxo-thiazolidin-4-one  
119. 5-[3-(1-Hydroxy-naphthalen-2-yl)-allylidene]-2-thioxo-thiazolidin-4-one  
120. 5-(2-Methyl-4-phenyl-pentylidene)-thiazolidine-2,4-dithione  
20 121. 5-[3-(4-Octadecyloxy-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one  
122. 5-(4-Diphenylamino-benzylidene)-2-thioxo-thiazolidin-4-one  
123. 5-(3,4,5-Trihydroxy-benzylidene)-thiazolidine-2,4-dithione  
124. 5-(4-Dimethylamino-2-methoxy-benzyl)-2-thioxo-thiazolidin-4-one  
125. 5-[3-(2-Benzoyloxy-4,5-dimethoxy-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one  
25 126. 5-[3-(2-Hydroxy-ethoxy)-benzylidene]-2-thioxo-thiazolidin-4-one  
127. 5-[2-(2-Hydroxy-ethoxy)-benzylidene]-thiazolidine-2,4-dithione  
128. 5-[4-(2-Hydroxy-ethoxy)-benzyl]-2-thioxo-thiazolidin-4-one  
129. Carboxylic acid *tert*-butyl ester 2-methoxy-4-[3-(4-oxo-2-thioxo-thiazolidin-5-ylidene)-propenyl]-phenyl ester  
30 130. 5-(3,5-Di-*tert*-butyl-2-hydroxy-benzylidene)-2-thioxo-thiazolidin-4-one  
131. 5-(2,4-Diethoxy-3-methyl-benzylidene)-thiazolidine-2,4-dithione  
132. 5-[3-(4-Methanesulphonyl-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one  
133. 5-(2-Hydroxy-5-methyl-benzylidene)-2-thioxo-thiazolidin-4-one  
134. 5-Benzo[*b*]thiophen-2-ylmethylene-thiazolidine-2,4-dithione  
35 135. 5-(5-Benzo[*b*]thiophen-2-yl-penta-2,4-dienylidene)-2-thioxo-thiazolidin-4-one

136. 5-(3-Naphthalen-2-yl-allylidene)-thiazolidine-2,4-dithione  
137. 2-Thioxo-5-[3-(2,6,6-trimethyl-cyclohex-1-enyl)-allylidene]-thiazolidin-4-one  
138. 5-(3-*tert*-Butyl-4-hydroxy-benzylidene)-2-thioxo-thiazolidin-4-one  
139. 5-(2,4-Bis-benzyloxy-benzylidene)-thiazolidine-2,4-dithione  
5 140. 5-(4-Benzyl-benzyl)-2-thioxo-thiazolidin-4-one  
141. 5-[3-(1*H*-Pyrrol-2-yl)-allylidene]-2-thioxo-thiazolidin-4-one  
142. 5-(5,6-Diethoxy-benzo[*b*]thiophen-2-ylmethylene)-thiazolidine-2,4-dithione  
143. 5-[3-(1-Methyl-1*H*-pyrrol-3-yl)-allylidene]-2-thioxo-thiazolidin-4-one  
144. 5-Cyclohexylmethylene-2-thioxo-thiazolidin-4-one  
10 145. 5-(2-Hydroxy-4,6-dimethoxy-benzylidene)-2-thioxo-thiazolidin-4-one  
146. 5-(4-Benzyloxy-2-hydroxy-benzylidene)-thiazolidine-2,4-dithione  
147. 5-(5-Benzyloxy-2-hydroxy-benzyl)-2-thioxo-thiazolidin-4-one  
148. 5-{3-[4-(Benzo[1,3]dioxol-5-ylmethoxy)-phenyl]-allylidene}-2-thioxo-  
thiazolidin-4-one  
15 149. 5-(4-Benzyloxy-3,5-dimethoxy-benzylidene)-2-thioxo-thiazolidin-4-one  
150. 5-(4-Benzyloxy-3,5-dihydroxy-benzylidene)-thiazolidine-2,4-dithione  
151. 5-(2,5-Bis-benzyloxy-benzyl)-2-thioxo-thiazolidin-4-one  
152. 5-[3-Methyl-5-(2,6,6-trimethyl-cyclohex-1-enyl)-penta-2,4-dienylidene]-  
thiazolidine-2,4-dithione  
20 153. 2-(2,4-Dithioxo-thiazolidin-5-ylidenemethyl)-benzoic acid  
154. 2-Methoxy-4-(4-oxo-2-thioxo-thiazolidin-5-ylmethyl)-phenyl acetate  
155. 2-Hydroxy-5-(4-oxo-2-thioxo-thiazolidin-5-ylidenemethyl)-benzoic acid  
156. 4-(2,4-Dithioxo-thiazolidin-5-ylidenemethyl)-benzoic acid  
157. 3-[4-(4-Oxo-2-thioxo-thiazolidin-5-ylmethyl)-phenyl]-acrylic acid  
25 158. 3-(4-Oxo-2-thioxo-thiazolidin-5-ylidenemethyl)-phenyl acetate  
159. [4-(2,4-Dithioxo-thiazolidin-5-ylidenemethyl)-phenoxy]-acetic acid  
160. 3-(4-Oxo-2-thioxo-thiazolidin-5-ylmethyl)-benzoic acid  
161. 5-(5,7-Dimethyl-4-oxo-4*H*-chromen-3-ylmethylene)-2-thioxo-thiazolidin-4-one  
162. 11-(2,4-Dithioxo-thiazolidin-5-ylidenemethyl)-1,4-dihydroxy-10-methoxy-5,8-  
30 dimethyl-1*H*-benzo[*e*]furo[3',4':3,4]benzo[*b*][1,4]dioxepine-3,7-dione  
163. 8-(4-Oxo-2-thioxo-thiazolidin-5-ylmethyl)-naphthalene-1-carboxylic acid  
164. 2-Acetoxy-5-(4-oxo-2-thioxo-thiazolidin-5-ylidenemethyl)-phenyl acetate  
165. 2-Amino-3-(2,4-dithioxo-thiazolidin-5-ylidenemethyl)-6,7-dimethyl-chromen-4-  
one  
35 166. 5-(6-Ethyl-4-oxo-4*H*-chromen-3-ylmethyl)-2-thioxo-thiazolidin-4-one



167. 5-(6,8-Dimethyl-4-oxo-4*H*-chromen-3-ylmethylene)-2-thioxo-thiazolidin-4-one  
168. Methyl 2-(2,4-dithioxo-thiazolidin-5-ylidenemethyl)-benzoate  
169. Methyl 3-(4-oxo-2-thioxo-thiazolidin-5-ylmethyl)-1*H*-indole-6-carboxylate  
170. 5-(1-*p*-Tolyl-ethylidene)-thiazolidine-2,4-dithione  
5 171. 5-[1-(4-Methoxy-phenyl)-ethyl]-2-thioxo-thiazolidin-4-one  
172. 5-[1-(3,5-Dihydroxy-phenyl)-ethylidene]-thiazolidine-2,4-dithione  
173. 2,6-Diacetoxy-4-(4-oxo-2-thioxo-thiazolidin-5-ylmethyl)-phenyl acetate  
174. 5-(3-Cyclohexyl-allylidene)-2-thioxo-thiazolidin-4-one  
175. 5-[5-(3,4-Diethoxy-2,5-dimethyl-phenyl)-penta-2,4-dienylidene]-2-thioxo-  
10 thiazolidin-4-one  
176. 2-Hydroxy-5-[3-(4-oxo-2-thioxo-thiazolidin-5-ylidene)-propenyl]-benzoic acid  
177. 3-[3-(4-Oxo-2-thioxo-thiazolidin-5-ylidene)-propenyl]-phenyl acetate  
178. 5-[3-(5,7-Dimethyl-4-oxo-4*H*-chromen-3-yl)-allylidene]-2-thioxo-thiazolidin-4-  
one  
15 179. 2-Acetoxy-5-[3-(4-oxo-2-thioxo-thiazolidin-5-ylidene)-propenyl]-phenyl acetate  
180. 5-[3-(6,8-Dimethyl-4-oxo-4*H*-chromen-3-yl)-allylidene]-2-thioxo-thiazolidin-4-  
one  
181. 5-(3-Phenyl-but-2-enylidene)-2-thioxo-thiazolidin-4-one  
182. 5-(3-Thiophen-2-yl-but-2-enylidene)-2-thioxo-thiazolidin-4-one  
20 183. 5-(2,4-Dimethoxy-benzyl)-thiazolidin-4-one  
184. 5-(2-Hydroxy-benzyl)-thiazolidin-4-one  
185. 5-(4-Diethylamino-2-hydroxy-benzyl)-thiazolidin-4-one  
186. 5-(2-Methyl-benzyl)-thiazolidin-4-one  
187. 5-[3-(4-Methoxy-phenoxy)-benzyl]-thiazolidin-4-one  
25 188. 5-(3,4,5-Trimethoxy-benzyl)-thiazolidin-4-one  
189. 5-(3-Hydroxy-4-methoxy-benzyl)-thiazolidin-4-one  
190. 5-(4-Diethylamino-benzyl)-thiazolidin-4-one  
191. 5-(4-Ethoxy-benzyl)-thiazolidin-4-one  
192. 5-(4-Methoxy-naphthalen-1-ylmethyl)-thiazolidin-4-one  
30 193. 5-(5-Methoxy-1*H*-indol-3-ylmethyl)-thiazolidin-4-one  
194. 5-(4-Hydroxy-3,5-dimethoxy-benzyl)-thiazolidin-4-one  
195. 5-(4-Isopropyl-benzyl)-thiazolidin-4-one  
196. 5-(2-Methyl-3-phenyl-allyl)-thiazolidin-4-one  
197. 5-(2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-thiazolidin-4-one  
35 198. 5-(2-Methyl-1*H*-indol-3-ylmethyl)-thiazolidin-4-one

199. 5-Benzofuran-2-ylmethyl-thiazolidin-4-one  
200. 5-(4-Hexyl-benzyl)-thiazolidin-4-one  
201. 5-(3-Amino-benzyl)-thiazolidin-4-one  
202. 5-(3,5-Dimethyl-1-phenyl-1*H*-pyrazol-4-ylmethyl)-thiazolidin-4-one  
5 203. 5-(2,2-Dimethyl-chroman-6-ylmethyl)-thiazolidin-4-one  
204. 5-(4-Dibutylamino-benzyl)-thiazolidin-4-one  
205. 5-(1-Hydroxy-naphthalen-2-ylmethyl)-thiazolidin-4-one  
206. 5-(4-Octadecyloxy-benzyl)-thiazolidin-4-one  
207. 5-(4-Methanesulphonyl-benzyl)-thiazolidin-4-one  
10 208. 5-(2,6,6-Trimethyl-cyclohex-1-enylmethyl)-thiazolidin-4-one  
209. 5-(1*H*-Pyrrol-2-ylmethyl)-thiazolidin-4-one  
210. 5-(1-Methyl-1*H*-pyrrol-3-ylmethyl)-thiazolidin-4-one  
211. 5-[4-(Benzo[1,3]dioxol-5-ylmethoxy)-benzyl]-thiazolidin-4-one  
  
15 211. 2-Hydroxy-5-(4-oxo-thiazolidin-5-ylmethyl)-benzoic acid  
212. 2-Hydroxy-5-(4-oxo-thiazolidin-5-ylmethyl)-benzoic acid  
213. 5-(5,7-Dimethyl-4-oxo-4*H*-chromen-3-ylmethyl)-thiazolidin-4-one  
214. 5-(6,8-Dimethyl-4-oxo-4*H*-chromen-3-ylmethyl)-thiazolidin-4-one  
215. 5-(1-Phenyl-ethyl)-thiazolidin-4-one  
20 216. 5-(1-Thiophen-2-yl-ethyl)-thiazolidin-4-one

The following Examples show some process variants which can be used for the synthesis of the compounds in accordance with the invention. However, they are not intended to be a limitation of the object of the invention. The structure of the compounds was  
25 proven by <sup>1</sup>H- and, where necessary, by <sup>13</sup>C-NMR spectroscopy. The purity of the substances was determined by C, H, N, P analysis as well as by thin-layer chromatography.

#### Example 1

##### 30 General Process A:

A solution of 5 mmol of aldehyde R-CHO, wherein R has the given significance, or of the corresponding ketone and 5 mmol of 2-thioxo-thiazolidin-4-one in 30 ml of abs. toluene is treated with catalytic amounts of piperidinium acetate and heated at reflux

for 5 to 10 hours. Thereafter, the mixture is cooled to 0°C. The precipitate is filtered off under suction, rinsed with diethyl ether and dried.

5-(4-Bromo-benzylidene)-2-thioxo-thiazolidin-4-one (1)

5 M.p. 226-7°C

5-Naphthalen-2-ylmethylene-2-thioxo-thiazolidin-4-one (2)

Orange-red crystals; m.p. 268-70°C

10 5-Thiophen-3-ylmethylene-2-thioxo-thiazolidin-4-one (3)

M.p.. 204°C (dec.)

5-(4-Hydroxy-benzylidene)-2-thioxo-thiazolidin-4-one (4)

Yellow crystals; m.p.. 214-6°C

15

5-(3,4-Diethoxy-benzylidene)-2-thioxo-thiazolidin-4-one (5)

Yellow-orange crystals; m.p. 186-7°C

5-[3-(5,6-Diethoxy-benzo[b]thiophen-2-yl)-allylidene]-2-thioxo-thiazolidin-4-one (6)

20 Brown crystals; m.p. 268°C

5-Thiophen-2-ylmethylene-2-thioxo-thiazolidin-4-one (7)

Yellow crystals; m.p. 223-5°C

25 5-Furan-2-ylmethylene-2-thioxo-thiazolidin-4-one (8)

Orange crystals; m.p. 231-33°C

5-[3-(3,4-Diethoxy-2,5-dimethyl-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one (9)

Brown crystals; m.p. 205-10°C

30

5-[1-(4-Chloro-phenyl)-ethylidene]-2-thioxo-thiazolidin-4-one (10)

Yellow crystals; m.p. 196-8°C

5-Pyridin-2-ylmethylene-2-thioxo-thiazolidin-4-one (11)

35 Olive green crystals; m.p. 250-5°C

5-(1-Phenyl-ethylidene)-2-thioxo-thiazolidin-4-one (12)

Yellow crystals; m.p. 166-8°C

5

5-(1-Thiophen-2-yl-ethylidene)-2-thioxo-thiazolidin-4-one (13)

Orange crystals; m.p. 218-20°C

5-(2-Hydroxy-benzylidene)-2-thioxo-thiazolidin-4-one (14)

10 M.p. 218°C (dec.)

5-(3,4-Dimethoxy-benzylidene)-2-thioxo-thiazolidin-4-one (15)

M.p. 187-9°C

15 5-(4-Isopropyl-benzylidene)-2-thioxo-thiazolidin-4-one (16)

M.p. 146-8°C

5-Naphthalen-1-ylmethylene-2-thioxo-thiazolidin-4-one (17)

M.p. 220-2°C

20

5-(5-Methyl-furan-2-ylmethylene)-2-thioxo-thiazolidin-4-one (18)

M.p. 227°C (dec.)

5-(4-Methoxy-benzylidene)-2-thioxo-thiazolidin-4-one (19)

25 M.p. 206°C (dec.)

5-(4-Ethoxy-benzylidene)-2-thioxo-thiazolidin-4-one (20)

M.p. 187-9°C

30 5-[3-(3,5-Di-tert-butyl-4-hydroxy-phenyl)-allylidene]-2-thioxo-thiazolidin-4-one (21)

Orange crystals; m.p. 205-10°C

5-(3-Benzo[b]thiophen-2-yl-allylidene)-2-thioxo-thiazolidin-4-one (22)

Orange crystals; m.p. 250°C

35

5-(3-Thiophen-2-yl-allylidene)-2-thioxo-thiazolidin-4-one (23)

Red-brown crystals; m.p. 213-6°C

5-(3-Naphthalen-2-yl-allylidene)-2-thioxo-thiazolidin-4-one (24)

5 Orange crystals; m.p. 256-8°C

5-[1-Methyl-3-(2,6,6-trimethyl-cyclohex-1-enyl)-allylidene]-2-thioxo-thiazolidin-4-one (25)

Yellow crystals; m.p. 189-10°C

10

5-(2-[1,3]Dioxolan-2-yl-6-fluoro-benzylidene)-2-thioxo-thiazolidin-4-one (26)

Beige crystals; m.p. 188-9°C

2-Thioxo-5-(2,6,6-trimethyl-cyclohex-1-enylmethylen)-thiazolidin-4-one (27)

15 Yellow crystals; m.p. 129-30°C

5-(4-Benzyl-benzylidene)-2-thioxo-thiazolidin-4-one (28)

Yellow-orange crystals; m.p. 210°C

20 5-(5,6-Diethoxy-benzo[b]thiophen-2-ylmethylen)-2-thioxo-thiazolidin-4-one (29)

Orange crystals; m.p. >250°C

5-[5-(5,6-Diethoxy-benzo[b]thiophen-2-yl)-penta-2,4-dienylidene]-2-thioxo-thiazolidin-4-one (30)

25 Dark brown crystals; m.p. 235-7°C

2-Thioxo-5-(1-p-tolyl-ethylidene)-thiazolidin-4-one (31)

Yellow crystals; m.p. 170-2°C

30 5-[1-(4-Methoxy-phenyl)-ethylidene]-2-thioxo-thiazolidin-4-one (32)

Yellow crystals; m.p. 164-6°C

5-[1-(3,4-Dichloro-phenyl)-ethylidene]-2-thioxo-thiazolidin-4-one (33)

Yellow crystals; m.p. 140-2°C

35

4-[4-(4-Oxo-2-thioxo-thiazolidin-5-ylidenemethyl)-benzyloxy]-benzonitrile (34)

Orange-brown crystals; m.p. 249-52°C

4-[4-(4-Oxo-2-thioxo-thiazolidin-5-ylidenemethyl)-phenoxy]-butyric acid (35)

5 Orange-brown crystals; m.p. 201-2°C

5-(11-Oxo-6,11-dihydro-dibenzo[b,e]oxepin-3-ylmethylene)-2-thioxo-thiazolidin-4-one (36)

Brown crystals; m.p. 270-2°C

10

5-(1H-Imidazol-2-ylmethylene)-2-thioxo-thiazolidin-4-one (37)

Orange-red crystals; m.p. 256°C

5-Benzo[b]thiophen-2-ylmethylene-2-thioxo-thiazolidin-4-one (38)

15 Yellow-orange crystals; m.p. 277-80°C

5-[3-Methyl-5-(2,6,6-trimethyl-cyclohex-1-enyl)-penta-2,4-dienylidene]-2-thioxo-thiazolidin-4-one (39)

Red crystals; m.p. 170-3°C

20

5-(3,5-Di-tert-butyl-4-hydroxy-benzyliden)-2-thioxo-thiazolidin-4-one (40)

Yellow crystals; m.p. 244-6°C

5-Benzylidene-2-thioxo-thiazolidin-4-one (41)

25 Yellow crystals; m.p. 202°C

5-(1H-Pyrrol-2-ylmethylene)-2-thioxo-thiazolidin-4-one (42)

LSM-0042541 BM 17.0564 17 AF 0090/1

Orange-red crystals; m.p. 272-4°C

30

5-(1-Methyl-1H-pyrrol-2-ylmethylene)-2-thioxo-thiazolidin-4-one (43)

Red-brown crystals; m.p. 248-50°C

Ethyl 2-(4-oxo-2-thioxo-thiazolidin-5-ylidenemethyl)-pyrrole-1-carboxylate (44)

35 Yellow crystals; m.p. 210-11°C

5-(4-Chloro-benzylidene)-2-thioxo-thiazolidin-4-one 5)

Yellow-orange crystals; m.p. 223-4°C

5 5-(3,4-Dichloro-benzylidene)-2-thioxo-thiazolidin-4-one (46)

Yellow-orange crystals; m.p. 234-5°C

### Example 2

10 General Process B:

1.6 mmol of 2,6-dimethyl-1,4-dihydro-3,5-pyridinedicarboxylic acid ethyl ester are added to a suspension of 1.2 mmol of 2-thioxo-thiazolidin-4-one derivative (Example 1) in 20 ml of toluene. The mixture is heated to 80°C for 22 hours, then the solution is  
15 filtered while warm. The residue is rinsed with ethyl acetate. The combined org. phases are concentrated, taken up in ethyl acetate and extracted with 1M HCl, dried over sodium sulphate and concentrated.

### Example 3

20 General Process C:

5 mmol of zinc dust in glacial acetic acid (5 ml/g zinc) are added to 1 mmol of rhodanine derivative (Example 2) divided into five portions in 30-60 minutes. Thereafter, the mixture is boiled at reflux for 2 to 24 hours. It is cooled to RT, infusorial  
25 earth is added and filtered off. The filtrate is treated with aqueous HCl and extracted with ethyl acetate. The combined org. phases are dried over sodium sulphate and concentrated. The residue is purified by chromatography (silica gel) with ethyl acetate/heptane.

30 Example 4

General Process D:

1 mmol of rhodanine derivative (Example 1) is dissolved in 40 ml of dioxan, treated with 1 mmol of P<sub>2</sub>S<sub>5</sub> and heated at reflux. After 2 to 10 hours the mixture is treated with  
35 active charcoal and filtered. The dioxan is removed under a vacuum and the residue is

crystallized with ethanol. For purification, it is treated with cold dimethylformamide, treated with active charcoal and precipitated with water.

#### General Process E:

5

10 mmol of thiazolidine-2,4-dione (Chem. Heterocycl. Compds. EN 2\_267-70, 1966) are stirred with 10 mmol of RCHO, in which R has the given significance, in 20 ml of methanol at room temperature for 60 min. The precipitate is filtered off under suction and recrystallized.

10

5-Naphthalen-1-ylmethylene-thiazolidine-2,4-dithione (47)

Red-brown crystals; m.p. 203°C (dec.)

5-Benzo[1,3]dioxol-5-ylmethylene-thiazolidine-2,4-dithione (48)

15

Red-brown crystals; m.p. 232°C (dec.)

5-(3-Benzo[b]thiophen-2-yl-allylidene)-thiazolidine-2,4-dithione (49)

Black crystals; m.p. 202-3°C

20

#### Example 5

Compounds of general formula (I) are investigated in a suitable assay for the capability of stimulating cyclic adenylyate cyclase.

25




Table I:

Example No.	Name	% cAMP (Test conc. 50µM)
<u>6</u>	5-[3-(5,6-Diethoxy-benzo[b]thiophen-2-yl)-allylidene]-2-thioxo-thiazolidin-4-one	8
<u>24</u>	5-(3-Naphthalen-2-yl-allyliden)-2-thioxo-thiazolidin-4-on	8
<u>25</u>	5-[1-Methyl-3-(2,6,6-trimethyl-cyclohex-1-enyl)-allylidene]-2-thioxo-thiazolidin-4-one	8
<u>27</u>	2-Thioxo-5-(2,6,6-trimethyl-cyclohex-1-enylmethylene)-thiazolidin-4-one	10
<u>39</u>	5-[3-Methyl-5-(2,6,6-trimethyl-cyclohex-1-enyl)-penta-2,4-dienylidene]-2-thioxo-thiazolidin-4-one	8
<u>40</u>	5-(3,5-Di-tert-butyl-4-hydroxy-benzylidene)-2-thioxo-thiazolidin-4-one	15
<u>49</u>	5-(3-Benzo[b]thiophen-2-yl-allylidene)-thiazolidine-2,4-dithione	10

## Patent Claims

1. Use of compounds of general formula (I)



5


in which

- m signifies a number between 0 and 8,  
 q signifies a number between 0 and 8  
 10 X signifies the group CH<sub>2</sub> or C=S, whereby A signifies a single bond and m signifies 0 when X signifies CH<sub>2</sub>,  
 A signifies a single or double bond  
 R<sub>1</sub>, R<sub>2</sub> signify hydrogen or lower alkyl, whereby R<sub>1</sub> and R<sub>2</sub> can be the same or different  
 and, when m signifies 2-8, R<sub>1</sub> and R<sub>2</sub> in the group CR<sub>1</sub>=CR<sub>2</sub> can have various  
 15 significances within the following sequence  
 R<sub>3</sub> signifies hydrogen or lower alkyl  
 Z signifies oxygen, sulphur  
 W signifies an optionally mono- or polysubstituted saturated or unsaturated  
 mono-, bi- or tricycle which can contain one or more hetero atoms,

20

for the preparation of medicaments for the treatment and prevention of metabolic bone disorders.

- 25 2. Compounds of general formula (I)



in which

m signifies a number between 0 and 8,

q signifies a number between 0 and 8

X signifies the group  $\text{CH}_2$  or  $\text{C}=\text{S}$ , whereby A signifies a single bond and m

5 signifies 0 when X signifies  $\text{CH}_2$ ,

A signifies a single or double bond

$\text{R}_1, \text{R}_2$  signify hydrogen or lower alkyl, whereby  $\text{R}_1$  and  $\text{R}_2$  can be the same or different and, when m signifies 2-8,  $\text{R}_1$  and  $\text{R}_2$  in the group  $\text{CR}_1=\text{CR}_2$  can have various significances within the following sequence

10  $\text{R}_3$  signifies hydrogen or lower alkyl

Z signifies oxygen, sulphur

W signifies an optionally mono- or polysubstituted saturated or unsaturated mono-, bi- or tricycle which can contain one or more hetero atoms,

15 whereas W is not phenyl, naphthyl, indolyl and thienyl, if X is  $\text{C}=\text{S}$ , Z is oxygen and m and q are both 0,

whereas W is not phenyl, furyl, thienyl and pyrrolyl, if X is  $\text{C}=\text{S}$ , Z is oxygen, A is a double bond and m is 0 or 1 and q is unequal 0 or m is unequal 0 and q is 2,

20

whereas W is not indolyl, if X is  $\text{C}=\text{S}$ , Z is oxygen, A is a double bond and m is 1 and 1 is 0,

whereas W is not 4-(2,5-di-tert. butyl-phenol), if X is methylene,

25 as well as their physiologically compatible salts, esters, optically active forms, racemates, tautomers, as well as derivatives which can be metabolized *in vivo* to compounds of general formula (I).

30 3. Medicament containing at least one compound of general formula (I) accordingly to claim 2 in admixture with usual pharmaceutical adjuvants and carrier materials

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/07250

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D277/36 C07D277/34 A61K31/425 C07D417/06 C07D417/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 783 888 A (SANKYO COMPANY LIMITED) 16 July 1997 (1997-07-16) the whole document	1-3
X	EP 0 677 517 A (ELI LILLY AND COMPANY) 18 October 1995 (1995-10-18) cited in the application claims	1,2
X	EP 0 604 983 A (MITSUBISHI KASEI CORPORATION) 6 July 1994 (1994-07-06) claims	1,2
X	EP 0 587 377 A (ELI LILLY AND COMPANY) 16 March 1994 (1994-03-16) claims	1,2
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

6 January 2000

Date of mailing of the international search report

14/01/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Henry, J

# INTERNATIONAL SEARCH REPORT

Interr. Natl Application No

PCT/EP 99/07250

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 434 394 A (ELI LILLY AND COMPANY) 26 June 1991 (1991-06-26) claims ---	1,2
X	EP 0 398 179 A (NISSHIN FLOUR MILLING CO) 22 November 1990 (1990-11-22) cited in the application claims ---	1,2
X	EP 0 391 644 A (ELI LILLY AND COMPANY) 10 October 1990 (1990-10-10) claims ---	1,2
X	EP 0 343 643 A (WARNER- LAMBERT COMPANY) 29 November 1989 (1989-11-29) claims ---	1,2
X	EP 0 316 790 A (NISSHIN FLOUR MILLING CO) 24 May 1989 (1989-05-24) claims ---	1,2
X	EP 0 237 138 A (YAMANOUCI PHARMACEUTICAL CO LTD) 16 September 1987 (1987-09-16) claims ---	1,2
X	EP 0 211 670 A (ELI LILLY AND COMPANY) 25 February 1987 (1987-02-25) cited in the application claims ---	1,2
X	WO 98 01445 A (SMITHKLINE BEECHAM S.P.A.) 15 January 1998 (1998-01-15) cited in the application claims ---	1-3
X	WO 96 26207 A (NISSAN CHEMICAL INDUSTRIES LTD) 29 August 1996 (1996-08-29) cited in the application claims ---	1,2
X	FR 2 196 797 A (ARIES ROBERT) 22 March 1974 (1974-03-22) the whole document ---	1,2
X	US 5 747 517 A (PANETTA JILL A.) 5 May 1998 (1998-05-05) claims ---	1,2
	-/--	

# INTERNATIONAL SEARCH REPORT

Intern. Patent Application No  
PCT/EP 99/07250

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HANS BEHRINGER ET AL: "Substituierte 5-methylen-rhodanine aus 5-chlormethylen-rhodaninen" CHEMISCHE BERICHTE., vol. 91, 1958, pages 2773-2782, XP002093301 WEINHEIM DE *pages 2773,2774,2779 ---	1
X	P.M. CHAKRABARTI ET AL: "An improved synthesis of substituted benzo[b]thiophen-2-carboxylic acids and related acids" TETRAHEDRON., vol. 25, 1969, pages 2781-2785, XP002093302 OXFORD GB page 2782 -page 2783 ---	1
X	CHEMICAL ABSTRACTS, vol. 120, no. 17, 25 April 1994 (1994-04-25) Columbus, Ohio, US; abstract no. 208602b, YASUHIRO O: page 101; XP002093303 abstract & JP 05 306224 A (WAKAMOTO PHARMA CO.LTD) 19 November 1993 (1993-11-19) ---	1,2
A	EP 0 691 129 A (ELI LILLY AND COMPANY) 10 January 1996 (1996-01-10) claims -----	1-3

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 99/ 07250

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 2 PARTIALLY  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 2 PARTIALLY

The initial phase of the search revealed a very large number of documents relevant to the issue of novelty. So many documents were retrieved that it is impossible to determine which parts of claim 2 (compounds per se) may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). Consequently, the search report with regard to said claim has been limited to a selection of retrieved novelty-affecting documents with special emphasis to the compounds illustrated by the examples and the list of preferred compounds of pages 10-16.

It should however be noted that the search and the search report can be considered as covering all claimed compounds of the prior art insofar those display an activity for the treatment and the prevention of metabolic bone disorders



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/07250

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0783888 A	16-07-1997	AU 7650396 A	03-07-1997
		CA 2193751 A	27-06-1997
		CZ 9603827 A	16-07-1997
		HU 9603607 A	28-07-1998
		JP 9235229 A	09-09-1997
		NO 965563 A	27-06-1997
		US 5804590 A	08-09-1998
EP 0677517 A	18-10-1995	CA 2144385 A	17-09-1995
		JP 7258235 A	09-10-1995
		US 5747517 A	05-05-1998
EP 0604983 A	06-07-1994	AT 145400 T	15-12-1996
		CA 2112331 A	29-06-1994
		DE 69306094 D	02-01-1997
		DE 69306094 T	03-04-1997
		DK 604983 T	09-12-1996
		ES 2097431 T	01-04-1997
		GR 3021746 T	28-02-1997
		JP 2845743 B	13-01-1999
		JP 6247945 A	06-09-1994
		US 5594016 A	14-01-1997
EP 0587377 A	16-03-1994	AU 676843 B	27-03-1997
		AU 4621893 A	17-03-1994
		CA 2105598 A	11-03-1994
		CN 1091006 A	24-08-1994
		CZ 9301814 A	16-03-1994
		EP 0915090 A	12-05-1999
		FI 933946 A	11-03-1994
		HU 70184 A	28-09-1995
		IL 106877 A	10-03-1998
		IL 119119 A	16-08-1998
		JP 6192091 A	12-07-1994
		MX 9305444 A	31-05-1994
		NO 933198 A	11-03-1994
		NO 981911 A	11-03-1994
		NZ 248573 A	27-02-1996
		PL 300335 A	21-03-1994
		US 5523314 A	04-06-1996
		US 5716975 A	10-02-1998
		US 5661168 A	26-08-1997
		ZA 9306492 A	02-03-1995
EP 0434394 A	26-06-1991	AT 169294 T	15-08-1998
		AU 639734 B	05-08-1993
		AU 6826690 A	27-06-1991
		CA 2032330 A	22-06-1991
		CN 1052668 A, B	03-07-1991
		DE 69032537 D	10-09-1998
		DE 69032537 T	21-01-1999
		ES 2121748 T	16-12-1998
		FI 906273 A	22-06-1991
		HU 216732 B	30-08-1999
		IL 96654 A	29-06-1995
		IL 108962 A	05-12-1996
		JP 4279573 A	05-10-1992
		MX 23803 A	28-02-1994

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter:      nal Application No

PCT/EP 99/07250

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0434394	A		NO 300458 B	02-06-1997
			NZ 236472 A	26-03-1993
			PT 96198 A,B	30-09-1991
			RU 2036915 C	09-06-1995
			RU 2050355 C	20-12-1995
			US 5387690 A	07-02-1995
			US 5216002 A	01-06-1993
EP 0398179	A	22-11-1990	DE 69024843 D	29-02-1996
			DE 69024843 T	30-05-1996
			KR 9608245 B	21-06-1996
			US 5116855 A	26-05-1992
			CA 2016665 A	19-11-1990
			JP 3072471 A	27-03-1991
EP 0391644	A	10-10-1990	AT 139531 T	15-07-1996
			AU 629322 B	01-10-1992
			AU 5293490 A	11-10-1990
			CA 2013599 A	07-10-1990
			DE 69027472 D	25-07-1996
			DE 69027472 T	05-12-1996
			DK 391644 T	15-07-1996
			ES 2088965 T	01-10-1996
			GR 3020500 T	31-10-1996
			JP 2290862 A	30-11-1990
			US 5356917 A	18-10-1994
			US 5691367 A	25-11-1997
EP 0343643	A	29-11-1989	AT 103175 T	15-04-1994
			AU 626863 B	13-08-1992
			AU 3505889 A	30-11-1989
			CA 1340247 A	15-12-1998
			DE 68914029 D	28-04-1994
			DE 68914029 T	07-07-1994
			DK 252089 A	26-11-1989
			EP 0565135 A	13-10-1993
			ES 2063073 T	01-01-1995
			FI 892522 A	26-11-1989
			IE 62214 B	11-01-1995
			JP 2062864 A	02-03-1990
			JP 2899309 B	02-06-1999
			KR 9702228 B	26-02-1997
			NO 892083 A	27-11-1989
			NZ 229266 A	23-12-1991
			PH 27092 A	26-02-1993
			PT 90662 A,B	30-11-1989
			US 5464856 A	07-11-1995
			US 5208250 A	04-05-1993
			US 5306822 A	26-04-1994
EP 0316790	A	24-05-1989	CA 1336837 A	29-08-1995
			DE 3883164 A	16-09-1993
			DE 3883164 T	02-12-1993
			ES 2059471 T	16-11-1994
			JP 1230565 A	14-09-1989
			JP 2645114 B	25-08-1997
			KR 9609424 B	19-07-1996
			US 4897406 A	30-01-1990

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/07250

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0237138	A	16-09-1987	AU 6740187 A DK 4487 A JP 63165368 A	09-07-1987 08-07-1987 08-07-1988
EP 0211670	A	25-02-1987	AT 52412 T AU 590312 B AU 6097286 A CA 1285572 A CN 1014891 B CY 1619 A DK 376986 A ES 2001075 A GR 862081 A HK 94791 A HU 42765 A IE 58718 B IL 79648 A JP 1902370 C JP 6025182 B JP 62042977 A KR 8700889 B LV 10866 A LV 10866 B MX 9203108 A NZ 217126 A PH 24517 A PT 83152 A, B SG 80391 G SU 1516012 A RU 2014329 C US 5356917 A US 5691367 A	15-05-1990 02-11-1989 12-02-1987 02-07-1991 27-11-1991 10-07-1992 10-02-1987 16-04-1988 24-12-1986 29-11-1991 28-08-1987 03-11-1993 12-12-1991 08-02-1995 06-04-1994 24-02-1987 02-05-1987 20-10-1995 20-04-1996 01-07-1992 27-01-1989 18-07-1990 01-09-1986 15-11-1991 15-10-1989 15-06-1994 18-10-1994 25-11-1997
WO 9801445	A	15-01-1998	EP 0912560 A US 5985905 A	06-05-1999 16-11-1999
WO 9626207	A	29-08-1996	AU 4731196 A JP 9235284 A ZA 9601478 A	11-09-1996 09-09-1997 28-08-1996
FR 2196797	A	22-03-1974	NONE	
US 5747517	A	05-05-1998	CA 2144385 A EP 0677517 A JP 7258235 A	17-09-1995 18-10-1995 09-10-1995
JP 05306224	A	19-11-1993	NONE	
EP 691129	A	10-01-1996	US 5476865 A CA 2153213 A JP 8040897 A	19-12-1995 07-01-1996 13-02-1996

**THIS PAGE BLANK (USPTO)**